

Silver-Haired Bats (*Lasionycteris noctivagans*) Found Ensnared on Burdock (*Arctium minus*)

Author(s): K. J. O. Norquay, A. K. Menzies, C. S. McKibbin, M. E. Timonin,
D. E. Baloun, and C. K. R. Willis

Source: Northwestern Naturalist, 91(3):339-342. 2010.

Published By: Society for Northwestern Vertebrate Biology

DOI: 10.1898/NWN10-08.1

URL: <http://www.bioone.org/doi/full/10.1898/NWN10-08.1>

BioOne (www.bioone.org) is an electronic aggregator of bioscience research content, and the online home to over 160 journals and books published by not-for-profit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Web site, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/page/terms_of_use.

Usage of BioOne content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

SILVER-HAIRED BATS (*Lasionycteris noctivagans*) FOUND ENSNARED ON BURDOCK (*Arctium minus*)

KJO NORQUAY, AK MENZIES, CS MCKIBBIN, ME TIMONIN, DE BALOUN, AND CKR WILLIS

Key words: *Arctium*, Burdock, *Lasionycteris noctivagans*, Silver-haired Bat

Burdock (*Arctium* spp.) is an invasive species from Eurasia, first reported in New England in 1638. It was common in Ontario by 1860, and is now considered an agricultural pest throughout its North American range (Royer and Dickinson 1999). Burdock's wide range can be attributed, in part, to the distinctive morphology of its seed casing, which has adhesive spines that readily attach to skin, fur, and feathers, enabling animal-based dispersal (Royer and Dickinson 1999). Burdock can pose a threat for small, flying vertebrates if the seed casing becomes entangled with the flight membrane of bats or the primary or secondary feathers of birds (for example, Lyon 1925; McNicholl 1988; Hager and others 2009). There are a number of published accounts of mortality for small birds found ensnared on Burdock including Ruby-throated Hummingbird (*Archilochus colubris*; McNicholl 1988; Nealen and Nealen 2000; Hinam and others 2004); Least Flycatcher (*Empidonax minimus*; Underwood and Underwood 2001; Hager and others 2009); Common Yellowthroat (*Geothlypis trichas*; Brown 1970); Blue-gray Gnatcatcher (*Poliophtila caerulea*; Brewer 1994; McNicholl 1994); Ruby-crowned Kinglet (*Regulus calendula*; Van Damme 2005); and Golden-crowned Kinglet (*Regulus satrapa*; Dawe 1974), among others (for example, Burton 1994; Dean 1994; McNicholl 1994; Underwood and Underwood 2001). These accounts suggest that mortality due to Burdock may be relatively common for small birds (3.5–15.2 g; Van Damme 2005), especially since many similar events likely go unreported. Although mortality caused by Burdock has been most widely reported for small birds, Burdock may pose a similar threat for small bats.

In 1892, approximately 12 bats, presumed to be Little Brown Bats (*Myotis lucifugus*) were found dead attached to Burdock in Illinois (Lyon 1925); Johnson (1933) reported a dead Eastern Red Bat (*Lasiurus borealis*) on Burdock in Virginia; Verts (1988) reported 2 dead Little Brown Bats attached to Burdock in Oregon; and Hendricks and others (2003) found 2 desiccated Western Long-eared Bats (*Myotis evotis*) on Burdock in Montana. In addition, unpublished observations of Silver-haired Bats (*Lasionycteris noctivagans*) ensnared on Burdock exist for Manitoba, Canada (SG Sealy and T Underwood, pers. comm.). Virtually all accounts were of dead animals except for a Big Brown Bat (*Eptesicus fuscus*) discovered alive in Illinois by Walley and others (1969); however, they did not report whether this bat survived or was released. Here we report the first published observations of Silver-haired Bats found attached to Burdock, and the first account of live bats found attached to Burdock that were successfully rehabilitated and released.

While radio-tracking Silver-haired Bats on the morning of 6 August 2009 in St. Leon, Manitoba, Canada (UTM: Zone14U, 529600E, 5467908N, NAD83), we found 2 juveniles without radio-transmitters tangled in the spines of seed heads approximately 1.5 m from the ground on 2 adjacent Burdock plants approximately 1.5 m apart. The bats were found in a small (0.02 km²) woodlot which, based on our radio-telemetry observations, contains a number of Silver-haired Bat roost trees (KJO Norquay and CKR Willis, unpubl. data). Both bats were stuck to the seed heads by their wing and interfemoral membranes, one facing head up and one head down. Each bat had accumulated approximately 20 fecal pellets in their interfemoral membranes, suggesting they were ensnared some-

time after foraging the night immediately previous to our discovery.

The bats were held in individual cloth bags and transported less than 0.5 km on foot to our field laboratory. Both bats were juvenile males, with masses of 8.9 and 10.6 g, respectively. The dry, wrinkled condition of the wing membranes suggested that the bats were dehydrated (Lollar and Schmidt-French 2002). In addition, the spines had caused between 5 and 10 small but visible holes (about 1 mm dia) in each wing membrane. Bats were initially lethargic but readily accepted water from a pipette and were hand-fed Mealworms (*Tenebrio molitor*). They were banded using numbered, lipped aluminium forearm bands (Porzana Ltd. East Sussex, UK), and placed in individual steel mesh bat cages in a quiet, dimly lit room for the rest of the day. Water was provided *ad libitum*. At dusk, the bats were provided additional water from a pipette and Mealworms. Prior to release, a small patch of fur ($<0.5 \text{ cm}^2$) was trimmed from the interscapular region of the larger bat, and a 0.53 g radio transmitter (LB-2NT, Holohil Systems Ltd., Carp, ON, Canada) was attached using a non-toxic, latex-based adhesive (Osto-bond, Montreal Ostomy, Vaudreuil, QC, Canada). Both bats were then hand-released at dusk within 20 m of the capture site, taking care to provide them a flight path which avoided the patch of Burdock.

We were able to confirm the survival and movement among roost trees of the radio-tagged individual for the next 4 d. We used a handheld telemetry receiver (R-1000, Communications Specialists Inc., Orange, CA, USA) and 5-element yagi antenna to track this individual each morning. The bat switched roost trees each night, using 3 roost trees before we could no longer detect a transmitter signal. All roost trees were located in the woodlot where we first found the bats. Given that the fall migration of Silver-haired Bats is likely underway in Southwestern Manitoba by early August (Barclay 1984), it is possible that the bat migrated when its signal disappeared.

It remains unclear how small bats and birds become caught in Burdock, but 2 hypotheses have been suggested. First, bats may be foraging for insects that live in or on Burdock or are caught on Burdock (Stager 1943). Bats might land on burrs while attempting to capture prey and become ensnared, or they become ensnared

while pursuing insects flying near Burdock. To our knowledge, there are no published observations of Silver-haired Bats gleaning prey from vegetation, but foraging Silver-haired Bats are known to specialize on swarming insects (Barclay 1985) and may become tangled if insects are highly concentrated near the seed heads of Burdock plants. Second, bats may collide with Burdock accidentally during normal flight through cluttered environments. Due to its position on Burdock, Johnson (1933) speculated that the Eastern Red Bat he observed was banking during a right-hand turn and became trapped accidentally. Bats have also been found caught on barbed wire (Johnson 1933), which is unlikely to attract insects, suggesting that foraging behavior alone does not explain the entrapment, and that accidental collision may be a more likely explanation. Juvenile bats are less able to maneuver than adults (Buchler 1980), which may increase their likelihood of collision with objects such as Burdock or barbed wire. Based on radio-tracking observations of other Silver-haired Bats in this woodlot, a roost tree is located within 3 m of the Burdock patch where we found the ensnared bats. The 2 bats we found were facing the roost tree which, along with evidence of recent feeding such as the presence of many fecal pellets, suggests that they were inexperienced juveniles that may have swooped too low when returning to their roost tree and thus became ensnared. The radio-tracked bat returned to that particular roost tree on the 2nd and 4th night following release.

The close proximity of the 2 bats we observed, caught in adjacent plants at approximately the same time, suggests that the 2 capture events might have been related. The calls of bats caught in Burdock may be attractive to conspecifics (Verts 1988), and it is possible that some bats are caught while investigating the vocalizations of other individuals which have already been ensnared. Recorded distress calls of bats are known to be attractive to conspecifics (Fenton and others 1976), and even other bat species (Russ and others 2004). In the case of our observations, once the 1st bat was caught, the 2nd bat may have investigated and been subsequently ensnared.

Mortality of bats due to collision or impalement with common objects in the environment

is not only limited to Burdock. Manville (1963) summarized reports of this kind of mortality in bats including impalement on fishhooks, barbed-wire fences, night-blooming Saguaro Cactus spines (*Carnegiea gigantea*), and rose-bush thorns, and entrapment of bats by other plants such as Spanish Moss (*Tillandsia usneoides*) and rosebushes. To our knowledge, however, perhaps with the exception of barbed-wire fences (DeBlase and Cope 1967, Iwen 1958, Long 1964), most of these examples represent highly isolated, rare occurrences. The widespread range of Burdock and the multiple reports in the literature of bat mortality for as many as 5 species suggest Burdock is a potentially serious, underestimated hazard for small flying vertebrates, including bats. We recommend that the objectives of future studies be to quantify the occurrence of bat and bird entrapment and mortality in areas heavily infested with Burdock, and to determine how and why birds and bats become ensnared. Most accounts of this phenomenon report that the animals failed to survive, so we were encouraged at our success in rehabilitating and releasing these individuals.

Acknowledgements.—We are grateful to property owners in the St. Leon area for site access. We thank T Underwood and S Sealy for sharing their observations and for help in finding relevant literature, and A Matheson for helping us find previous reports of vertebrates ensnared on Burdock. Funding was provided by a Natural Sciences and Engineering Research Council (NSERC, Canada) Discovery Grant and Sustainable Development Innovation Fund Grant (Manitoba Conservation) to CKRW, as well as NSERC Undergraduate Student Research Awards to AKM and CSM. P Hendricks, T Jung and an anonymous reviewer provided comments that improved the manuscript.

LITERATURE CITED

- BARCLAY RMR. 1984. Observations on the migration, ecology and behaviour of bats at Delta Marsh, Manitoba. *Canadian Field-Naturalist* 98:331–336.
- BARCLAY RMR. 1985. Long- versus short-range foraging strategies of hoary (*Lasiurus cinereus*) and silver-haired (*Lasionycteris noctivagans*) bats and the consequences for prey selection. *Canadian Journal of Zoology* 63:2507–2519.
- BREWER AD. 1994. Blue-gray gnatcatcher killed by entanglement on burdock. *Ontario Birds* 12:115–116.
- BROWN RD. 1970. Yellowthroat caught in common burdock. *The Wilson Bulletin* 82:464–465.
- BUCHLER ER. 1980. The development of flight, foraging, and echolocation in the little brown bat (*Myotis lucifugus*). *Behavioural Ecology and Sociobiology* 6:211–218.
- BURTON JF. 1994. Barn swallow trapped by greater burdock. *British Birds* 87:144.
- DAWE N. 1974. Golden-crowned kinglet death on burdock. *Murrelet* 55:29.
- DEAN T. 1994. Goldcrest corpse stuck on lesser burdock. *British Birds* 87:145.
- DEBLASE AF, COPE JB. 1967. An Indiana bat impaled on barbed wire. *American Midland Naturalist* 77:238.
- FENTON MB, BELLWOOD JJ, FULLARD JH, KUNZ TH. 1976. Responses of *Myotis lucifugus* (Chiroptera: Vespertilionidae) to calls of conspecifics and other sounds. *Canadian Journal of Zoology* 54:1443–1448.
- HAGER SB, DZIADYK B, MCKAY KJ. 2009. Bird-plant entanglement: A review and addition of the least flycatcher. *Wilson Journal of Ornithology* 121:648–651.
- HENDRICKS P, CARLSON J, CURRIER CJ. 2003. Fatal entanglement of western long-eared myotis in burdock. *Northwestern Naturalist* 84:44–45.
- HINAM HL, SEALY SG, UNDERWOOD TJ. 2004. Ruby-throated Hummingbird, *Archilochus colubris*, entanglements in burdock, *Arctium* spp., at Delta Marsh, Manitoba. *Canadian Field-Naturalist* 118:85–89.
- IWEN FA. 1958. Hoary bat the victim of a barbed wire fence. *Journal of Mammalogy* 39:438.
- JOHNSON PB. 1933. Accidents to bats. *Journal of Mammalogy* 14:156–157.
- LOLLAR AL, SCHMIDT-FRENCH BA. 2002. Captive care and medical reference for the rehabilitation of insectivorous bats, 2nd edition. Mineral Wells, TX: Bat World Publication. 340 p.
- LONG CA. 1964. Red Bat impaled on barbed wire. *Transactions of the Kansas Academy of Science* (1903) 67:201.
- LYON MW JR. 1925. Bats caught by burdocks. *Journal of Mammalogy* 6:280.
- MANVILLE RH. 1963. Accidental mortality in bats. *Mammalia* 27:361–366.
- MCNICHOLL MK. 1988. Bats and birds stuck on burdock. *Prairie Naturalist* 20:157–160.
- MCNICHOLL MK. 1994. Additional records of birds caught in burdock. *Ontario Birds* 12:117–119.
- NEALEN JH, NEALEN PM. 2000. Ruby-throated hummingbird death by common burdock. *Wilson Bulletin* 112:421–422.
- ROYER F, DICKINSON R. 1999. Weeds of Canada and the Northern United States: A guide to identification. Edmonton, AB: University of Alberta Press. 434 p.
- RUSS JM, JONES G, MACKIE IJ, RACEY PA. 2004. Interspecific responses to distress calls in bats

- (Chiroptera: Vespertilionidae): A function for convergence in call design? *Animal Behaviour* 67: 1005–1014.
- STAGER KE. 1943. California leaf-nosed bat trapped by desert shrub. *Journal of Mammalogy* 24:396.
- UNDERWOOD TJ, UNDERWOOD RM. 2001. Observations on burdock-killed birds in King's Park, Winnipeg, MB. *Blue Jay* 59:64–69.
- VAN DAMME LM. 2005. Burdock entrapment of a ruby-crowned kinglet in the Creston Valley. *Wildlife Afield* 2:14–15.
- VERTS BJ. 1988. Two bats caught on a plant. *Murrelet* 69:36–38.
- WALLEY HD, SOUTHERN WE, ZAR JH. 1969. Big brown bat entangled in burdock. *American Midland Naturalist* 82:630.
- Department of Biology and Centre for Forest Interdisciplinary Research, University of Winnipeg, Winnipeg, MB R3B 2E9 (KJON, AKM, CSM, MET, DEB, CKRW); corresponding author c. willis@uwinnipeg.ca. Submitted 1 March 2010, accepted 16 June 2010. Corresponding Editor: Thomas Jung.*